

# Indicators 2013



## Patient Safety Award

- Emergency readmission within 28 days of discharge following hip fracture (65+)
- Summary Hospital-level Mortality Index (SHMI)
- Risk adjusted mortality index
- Readmission rate within seven days of delivery
- Infection rate following caesarean section
- Rate of deaths in hospital within 30 days of non-elective surgery
- Rate of deaths in hospital within 30 days of elective surgery
- Rates of deaths in hospital within 30 days of emergency admission for hip fractures (FnoF; aged 65+)
- Rate of deaths in hospital within 30 days of emergency admission with a heart attack (MI: aged 35 to 74)
- Rate of deaths in hospital within 30 days of emergency admission for a stroke
- Rate of deaths in low mortality HRGs (HRG3.5)
- Rate of decubitus ulcer (for patients with length of stay over four days)
- Post-operative wound infection
- Complications of anaesthesia
- Foreign body left in during procedure
- Post-operative pulmonary embolism or deep vein thrombosis
- Post-operative sepsis
- Rate of accidental puncture or laceration during surgery
- Birth trauma - injury to neonate

## Quality of Care Award

- Day case conversion to inpatient rate (vs national rates, case mix adj per BADS)
- Patient reported outcomes score (across 4 procedures, per EQ5D)
- Rate of emergency readmission to hospital (>16; 28 days)
- Percentage of elective admissions where planned procedure not carried out (not patient decision)
- Summary Hospital-level Mortality Index (SHMI)
- Risk adjusted length of stay
- Risk adjusted mortality index
- Percentage of patients >65 with fractured neck of femur with pre-op LoS<=2
- Cancer patients seen within 2 weeks - all suspected cancers
- Discharge to usual place of residence within 56 days of emergency admission for patients with stroke
- Discharge to usual place of residence within 28 days of emergency admission for patients with a hip fracture (FnoF; aged 65+)
- Admitted patients' waiting time from point of referral to treatment

## Data Quality Award

- Depth of coding (not case mix adjusted)
- Percentage of coded episodes with signs and symptoms as a primary diagnosis
- Percentage of uncoded episodes

## 40Top Awards

Revised annually to take into account of newly-available performance information, this year's indicators include:

- Reported C-difficile rate for patients aged 65 and over
- Day case rate (relative weighted performance across BADS directory)
- Day case conversion to inpatient rate (vs national rates, case mix adj per BADS)
- Depth of coding (not case mix adjusted)
- Percentage of coded episodes with signs and symptoms as a primary diagnosis
- Percentage of uncoded episodes
- Inpatient survey (overall care question)
- Percentage of outpatient first appointments not attended (specialty adjusted)
- Rate of emergency readmission to hospital (>16; 28 days)
- Emergency readmission within 28 days of discharge following hip fracture (65+)
- Percentage of elective admissions where planned procedure not carried out (not patient decision)
- Reference Cost Index (RCI)
- Summary Hospital-level Mortality Index (SHMI)
- Staff survey (overall job satisfaction question)
- Risk adjusted length of stay
- Risk adjusted mortality index
- Rate of emergency readmission to hospital following AMI within 28 days
- Rate of emergency readmission to hospital within 14 days - COPD
- Percentage of elective inpatients admitted on day of procedure
- Patient misadventure rate (ICD-based)
- Percentage of patients >65 with fractured neck of femur with pre-op LoS <=2
- Unnecessary admissions via A&E (zero LoS as % emergency)

## Most Improved Hospital Award

Revised annually to take into account of newly-available performance information, this year's indicators include:

- Reported C-difficile rate for patients aged 65 and over
- Day case rate (relative weighted performance across BADS directory)
- Day case conversion to inpatient rate (vs national rates, case mix adj per BADS)
- Depth of coding (not case mix adjusted)
- Percentage of coded episodes with signs and symptoms as a primary diagnosis
- Percentage of uncoded episodes
- Inpatient survey (overall care question)
- Percentage of outpatient first appointments not attended (specialty adjusted)
- Patient reported outcomes score (across 4 procedures, per EQ5D)
- Rate of emergency readmission to hospital (>16; 28 days)
- Emergency readmission within 28 days of discharge following hip fracture (65+)
- Percentage of elective admissions where planned procedure not carried out (not patient decision)
- Reference Cost Index (RCI)
- Summary Hospital-level Mortality Index (SHMI)
- Staff survey (overall job satisfaction question)
- Risk adjusted length of stay

- Risk adjusted mortality index
- Rate of emergency readmission to hospital following AMI within 28 days
- Rate of emergency readmission to hospital within 14 days - COPD
- Percentage of elective inpatients admitted on day of procedure
- Patient misadventure rate (ICD-based)
- Percentage of patients >65 with fractured neck of femur with pre-op LoS <=2
- Unnecessary admissions via A&E (zero LoS as % emergency)